

Department of Water Government of Western Australia



Dathagnoorara Water Reserve Drinking Water Source Protection Plan

> REPORT NO. 71 June 2007

Water Resource Protection Series



Dathagnoorara Drinking Water Source Protection Plan

Carnamah and Coorow Town Water Supplies

Department of Water

Water Resource Protection Series

Report WRP 71

June 2007

Department of Water

Level 4, 168 St Georges Terrace Perth Western Australia 6000

<www.water.wa.gov.au>

Telephone +61-8-6364 7600 Facsimile +61-8-6364 7601

For more information about this report, contact

Program Manager Protection Planning, Water Source Protection Branch, or send an email to drinkingwater@water.wa.gov.au.

June 2007

ISSN 1326-7442

Subject of cover photograph

Dathagnoorara production bore 1/91, taken by Alana Thorpe

Contents

С	Contentsiii							
P	reface	۶۷	1					
S	umma	aryv	i					
1	1 Drinking water source overview1							
	1.1 1.2 1.3	Existing water supply system 1 Water treatment 1 Catchment details 2 1.3.1 Physiography 2 1.3.2 Climate 2 1.3.3 Hydrogeology 2	2222					
	1.4 1.5	Future water supply requirements 3 Protection and allocation 3 1.5.1 Existing water source protection 3 1.5.2 Current allocation licence 3	3					
2	Wate	er quality monitoring and contamination risks6	5					
	2.1 2.2 2.3 2.4 2.5	Microbiological contaminants 6 Health related characteristics 6 Aesthetic characteristics 7 Groundwater bores 7 Mineral exploration bores 8	5 7 7					
3	Land	use assessment9)					
	3.1 3.2	Existing land uses						
4	Catc	hment protection strategy11						
	4.1 4.2 4.3 4.4 4.5 4.6 4.7	Protection objectives 11 Proclaimed area 11 Priority classifications 11 Protection Zones 11 Land use planning 13 Best management practices 13 Surveillance and By-law enforcement 14	 					
	4.8 4.9	Emergency response						
5	Reco	ommendations)					
A	openc	dices21						
G	lossa	ry26	5					
R	eferer	nces and Recommended Reading29)					
С	ontrib	utors						

Appendices

Appendix A - Summary of Submissions	21
Appendix B- Water quality	22
Appendix C - Photographs	25

Figures

Figure 1 Dathagnoorara Water Reserve locality map	4
Figure 2 Dathagnoorara Water Reserve	
Figure 3 Land use and tenure within and surrounding the Dathagnoorara Water	
Reserve	10
Figure 4 Priority classifications and protection zones for Dathagnoorara Water	
Reserve	12

Tables

Table 1 Land use	. potential water	quality risks	s and recommended s	strategies15

Preface

The Department of Water has prepared this Drinking Water Source Protection Plan to report on the activities and risks to water quality within the Dathagnoorara Water Reserve, and to recommend management strategies to address these.

A safe drinking water supply is critical to the well-being of the community and catchment protection is necessary to help avoid, minimise or manage risks to water quality. The Department is committed to protecting drinking water sources to ensure the continued supply of 'safe, good quality drinking water' to consumers.

The Australian Drinking Water Guidelines recommend a multiple barrier risk based approach to protect public drinking water sources. Protection of drinking water catchments is the 'first barrier', with subsequent barriers implemented at the water storage, treatment and distribution stages of a water supply system. Catchment protection includes understanding the catchment, the hazards and hazardous events that can compromise drinking water quality, and developing and implementing preventive strategies and operational controls to ensure the safest possible water supply from our surface water dams and groundwater aquifers.

This plan details the location and boundary of the drinking water reserve, which provides potable water to the Carnamah and Coorow Town Water Supplies. It describes the water supply system, discusses existing and future usage of the water source, identifies risks and recommends management approaches to maximise protection of the catchment.

This plan should be used to guide State and Local Government land use planning decisions. It should be recognised in the Shire of Carnamah Town Planning Scheme, consistent with the Western Australian Planning Commission's Statement of Planning Policy No. 2.7 *Public Drinking Water Source Policy*. Other stakeholders should use this document as a guide for protecting the quality of water in the Dathagnoorara Water Reserve.

Stages in development of a Plan	Comment
Prepare Drinking Water Source	Prepared following a catchment survey and preliminary
Protection Assessment	information gathering.
Conduct stakeholder consultation	Advice sought from key stakeholders using the
	assessment as a tool for information and discussion.
Prepare Draft Drinking Water	Draft Plan developed taking into account input from
Source Protection Plan	stakeholders and any additional advice received.
Release Draft Drinking Water	Draft Plan released for a six week public consultation
Source Protection Plan	period.
Publish approved Drinking Water	Final Plan published after considering advice
Source Protection Plan	received in submissions. Includes recommendations
	on how to protect the catchment.

The stages involved in preparing a Drinking Water Source Protection Plan are:

Summary

Carnamah and Coorow are located in the Mid West Region of Western Australia, 307 and 280 kilometres north north east of Perth respectively. Both towns are supplied with water from two Water Corporation bores, located within the Dathagnoorara Water Reserve, approximately 20 kilometres south west of Carnamah. The Dathagnoorara Water Reserve was proclaimed under the *Country Areas Water Supply Act 1947* in 1992 to protect the public drinking water source.

The bores draw water from the Parmelia Formation, which was deposited in the Late Jurassic and consists of interbedded sandstone, claystone, siltstone and shale. This formation forms a semi-confined aquifer of regional extent.

The water table in the area of the Dathagnoorara Water Reserve is greater than 90 metres deep and the bores are screened at a depth of at least 150 metres. Combined with the semi-confined nature of the aquifer this results in a low risk of contamination to the drinking water source from surrounding agricultural land uses.

There is, however, a degree of risk to the drinking water source as the aquifer is not completely confined. Therefore, it is proposed that the boundary of the Dathagnoorara Water Reserve be amended to include the additional area formed by 300 metre Wellhead Protection Zones (WHPZ), in order to protect the area around the production bores where the water source is most vulnerable to contamination. The current gazetted water reserve will be managed for Priority 1 source protection while the remainder of the proposed Dathagnoorara Water Reserve will be managed for Priority 2 source protection. The boundary and priority classifications have been determined to provide the appropriate level of protection for the drinking water source, while recognising the rights of landowners to continue established approved land use activities.

The following strategies are recommended to protect the Dathagnoorara Water Reserve:

- extend the boundary of the Water Reserve to include the proposed WHPZ, in order to ensure the appropriate level of protection;
- the Water Reserve boundary, priority classifications and management principles outlined in this plan should be incorporated into the Shire of Carnamah's Town Planning Scheme and other applicable schemes and strategies;
- any development proposals that are inconsistent with the Water Quality Protection Note *Land use compatibility in Public Drinking Water Source Areas* should be referred to the Department of Water for comment; and

• best management practices for current or approved land uses in the Dathagnoorara Water Reserve should be implemented.

1 Drinking water source overview

1.1 Existing water supply system

Carnamah and Coorow are located in the Mid West Region of Western Australia approximately 307 and 280 kilometres north north east of Perth (see Figure 1). Both towns are service centres for local mining industries and broad acre agriculture, including cereal cropping, sheep and cattle grazing and wildflower farming. Carnamah and Coorow have populations of approximately 310 and 250 respectively.

Dathagnoorara Water Reserve is located within the Shire of Carnamah approximately 20 kilometres south west of the Carnamah townsite (see Figure 1). The Dathagnoorara wellfield consists of two production bores (1/91 and 2/86), approximately 120 metres apart and drilled into the Parmelia formation (see Figure 2 for bore locations). Bore 1/91 is screened between 152 and 181 metres with a static water level of 89 metres and bore 2/86 is screened between 150 and 200 metres with a static water level of 92 metres. Both bores are capped and the surface casing has been cemented down to 20 metres on bore 2/86 and to 17.4 metres on bore 1/91, which prevents any contamination of the water source in the sandy layers close to the surface.

The water from the production bores is treated and pumped into the Dathagnoorara collector tank, where it is disinfected and then pumped through the Dathagnoorara supply main to the Carnamah and Coorow reticulation and town storage tanks. Off-tanks from the main line also provide water for nearby farms. The wellfield currently supplies approximately 400 services.

1.2 Water treatment

Groundwater extracted from the Dathagnoorara bores is first aerated to raise the pH and reduce iron levels of the raw water. All water is then disinfected by chlorination prior to supply as drinking water.

It should be recognised that although treatment and disinfection are essential barriers used to ensure a safe, good quality, aesthetically acceptable drinking water supply, catchment management and water source protection are fundamental 'first barriers' for the protection of water quality. This approach is endorsed by the Australian Drinking Water Guidelines (ADWG) (NHMRC & NRMMC, 2004) and reflects a multiple barrier 'catchment to consumer' risk based approach for the provision of safe drinking water to consumers.

1.3 Catchment details

1.3.1 Physiography

Dathagnoorara Water Reserve is located on the Dandaragan Plateau, a sand and laterite capped plateau overlying Cretaceous sediments. The Plateau is flat to gently undulating, ranging in elevation from 140 to 260 metres above sea level. The most significant topographical feature in the area is the Dandaragan Scarp, which is located on the western boundary of the Plateau and rises up to around 50 metres above land further to the west.

Deep yellow sands are common in the uplands while the valleys are dominated by deep red and yellow brown sands.

1.3.2 Climate

Carnamah experiences a mild Mediterranean-type climate with hot, dry summers and cool, wet winters. Average rainfall is 363.8 millimetres, with the majority falling between May and August. Average daily maximum and minimum temperatures in January are 36.1℃ and 18.7 ℃ and in July are 17.1 ℃ and 7.7℃ respectively (Bureau of Meteorology, 2007).

1.3.3 Hydrogeology

Dathagnoorara wellfield is located to the west of the Urella Fault in the Dandaragan Trough, a major structural unit of the Perth Basin. The trough is bounded on the east by the Urella and Darling Faults and on the west by the Beagle Fault System.

The Parmelia Formation occurs between the Dandaragan Scarp and the Urella Fault and is separated from the underlying Yarragadee Formation by the Otorowiri Member, a siltstone aquiclude. The Parmelia Formation is of Late Jurassic age and consists of interbedded sandstone, claystone, siltstone and shale. The total thickness of the sandstone in the Parmelia Formation is variable but is generally in the order of 100 to 200 metres thick. The Yarragadee Formation is of Middle to Late Jurassic age and consists of sandstone, siltstone, shale and claystone up to 2500 metres thick (Water Authority, 1995).

The Dathagnoorara bores draw water from the Parmelia Formation. The Parmelia Formation forms an aquifer of regional extent, which has a complex flow system and contains large resources of fresh to brackish water. At its surface the aquifer is recharged by direct infiltration of rainfall and associated runoff, however, the presence of interbedded shale produces semi-confining layers beneath the Dathagnoorara borefield. The depth to the watertable in the area of the Dathagnoorara bores is more than 90 metres, and flow is in a general northerly direction, although within the area of the Dathagnoorara Water Reserve there also appears to be some leakage into the Yarragadee Formation below.

Carbon 14 dating techniques have been used to determine the average age of water in the Dathagnoorara bores. The average corrected age of the water in Bore 2/86 was found to be 6730 years. This provides a good indication of the considerable length of time it takes for water infiltrating at the recharge area to be withdrawn by the bores.

1.4 Future water supply requirements

The existing wellfield adequately meets the current requirements of Coorow and Carnamah. At this stage the Water Corporation considers that the current source should also be sufficient to meet future demand.

1.5 Protection and allocation

1.5.1 Existing water source protection

The Dathagnoorara Water Reserve was proclaimed in 1992 under the *Country Areas Water Supply (CAWS) Act 1947.* By-laws are applied under this Act to ensure protection of the water source against contamination. The gazetted Water Reserve covers Crown Reserve 25198 (see Figure 2).

1.5.2 Current allocation licence

Water resource use and conservation in Western Australia is administered by the Department of Water in accordance with the *Rights in Water and Irrigation (RIWI) Act 1914.* Under this Act, the right to use and control surface and groundwater is vested with the Crown. This Act requires licensing of groundwater abstraction within proclaimed groundwater areas.

The Dathagnoorara borefield is located within the Arrowsmith Groundwater Area. The resource is managed in accordance with this Department's Interim Sub-regional Allocation Strategy titled *Managing the Water Resources of the Arrowsmith Groundwater Area, WA* (Water and Rivers Commission, 2002). The Water Corporation hold a Groundwater Well Licence (no. 65345) to draw 400 000 kilolitres per annum from the Dathagnoorara wellfield for public water supply purposes. Current abstraction is lower than this, with a volume of 277 095 kilolitres abstracted during the 2005/06 licence period.




Figure 1 Dathagnoorara Water Reserve locality map




Figure 2 Dathagnoorara Water Reserve

2 Water quality monitoring and contamination risks

The Water Corporation regularly monitors the raw water quality from the Dathagnoorara wellfield for microbiological contamination, health related and aesthetic (non-health related) characteristics in accordance with the ADWG. The results of this monitoring are then reviewed by an intergovernmental committee, chaired by the Department of Health, called the Advisory Committee for the Purity of Water.

A water quality summary for the Dathagnoorara wellfield is presented in Appendix B. For more information on water quality, see the Water Corporation's most recent Drinking Water Quality Annual Report at <www.watercorporation.com.au> > Water > Water Quality > Downloads > access the most recent Annual Report.

Contamination risks relevant to the Dathagnoorara Water Reserve are described below.

2.1 Microbiological contaminants

Pathogens are types of micro-organisms that are capable of causing diseases. These include bacteria (such as *Escherichia coli*), protozoa (such as *Cryptosporidium* and *Giardia*) and viruses. In water supplies the pathogens of concern that can cause illness are mostly found in the faeces of humans and domestic animals. *Escherichia coli* counts are a way of measuring these pathogens and are an indicator of faecal contamination.

Pathogen contamination of a drinking water source is influenced by the existence of pathogen carriers (ie humans and domestic animals, such as dogs or cattle); their subsequent transfer to and movement in the water source; and the ability of the pathogen to survive in the water source.

The effects on people from consuming drinking water that is contaminated with pathogens is considerably varied, ranging from mild illness (such as stomach upset or diarrhoea) to death. This was the case in Walkerton, Canada in 2000, where seven people died due to the contamination of a pathogenic strain of *Escherichia coli* in the town water source and supply (NHMRC & NRMMC, 2004b). Preventing the introduction of pathogens into the water source is the most effective barrier in avoiding this public health risk.

2.2 Health related characteristics

A number of chemicals (organic and inorganic) are of concern in drinking water from a health perspective because they are potentially toxic to humans. Chemicals usually occur in drinking water sources attached to suspended material such as soil particles and may result from natural leaching from mineral deposits or from different land uses (NHMRC & NRMMC, 2004b)

Pesticides include agricultural chemicals such as insecticides, herbicides, nematicides, rodenticides and miticides. Contamination of a drinking water source by pesticides may occur as a result of accidental spills, incorrect or over use and leakage from storage areas. In such cases, prompt action is required to notify relevant authorities and clean up the spill.

Nutrients (such as nitrogen) can enter drinking water supplies from leaching of fertiliser, septic tanks, and from faeces of domestic animals (such as cattle grazing on the land). Nitrate and Nitrite (ions of Nitrogen) can be toxic to humans at high levels, with infants less than 3 months old being most susceptible (NHMRC & NRMMC, 2004).

Hydrocarbons (fuels, oils, solvents) are potentially toxic to humans, and harmful byproducts may be formed when they are combined with chlorine in water treatment processes. Hydrocarbons can occur in water supplies from pollution events from vehicle accidents, refuelling and leakage from storage areas.

2.3 Aesthetic characteristics

Impurities in drinking water can affect the aesthetic qualities of water such as appearance, taste, smell and 'feel'. Such impurities are not necessarily hazardous to human health, for example water that is cloudy and has a distinctive colour may not be harmful (NHMRC & NRMMC, 2004b).

Iron and dissolved organic matter can affect the colour and appearance of water, and salinity can affect the taste. The ADWG have set limits on water quality characteristics to meet aesthetic requirements of consumers.

Some properties such as pH can contribute to the corrosion and encrustation of pipes. The ADWG also sets out aesthetic guidelines for these types of water quality characteristics.

2.4 Groundwater bores

Under the provisions of sections 26D and 5C of the *RIWI Act*, a licence is required to construct a bore or extract water (unless exempt under the *RIWI* Exemption and Repeal (Section 26C) Order 2001) within a proclaimed groundwater area. The Dathagnoorara Water Reserve is located within the Arrowsmith Groundwater Area.

Any bores drilled near to a public drinking water supply bore have the potential to contaminate the drinking water source. For example, a poorly constructed bore may introduce contaminants through surface leakage down the outside of the bore casing into an otherwise uncontaminated aquifer. If a public drinking water source bore is nearby, it may abstract the contaminated water.

It is important to ensure that any bores are appropriately located and constructed in order to prevent contamination and other impacts on the public drinking water source. This will be assessed through the Department of Water's water licensing process where applicable under the *RIWI Act.*

All bores should be constructed in accordance with *Minimum Construction Requirements for Water Bores in Australia* (National Minimum Bore Specifications Committee, 2003).

2.5 Mineral exploration bores

Any drilling undertaken in close proximity to a drinking water source area has the potential to contaminate the source by providing a pathway for contaminants to travel from the ground surface down into the aquifer from which the production bores draw from. Where possible, exploration drilling should be kept outside of the Water Reserve and downstream of the production bores. All drilling undertaken in close proximity to the production bores must be appropriately backfilled and sealed in order to prevent contamination of the water source.

3 Land use assessment

3.1 Existing land uses

The current gazetted Dathagnoorara Water Reserve is Crown Land and is zoned for public purposes under the Shire of Carnamah Town Planning Scheme. Land surrounding the gazetted Water Reserve, which is included in the proposed Dathagnoorara Water Reserve is privately owned and zoned rural (see Figure 3). Broad acre cereal cropping including wheat, barley, lupins and canola is the primary agricultural land use, with some low intensity stock grazing (sheep and cattle) also occurring in the area.

3.2 Proposed land uses

Under the current rural zoning there is the potential for intensification of agricultural land uses in the area. Nutrient outputs are likely to be greater from more intensive agriculture, which could potentially result in a higher risk of contamination to the water source than the current extensive agriculture. Any proposals to intensify or change the agricultural land use that are inconsistent with the Water Quality Protection Note *Land use compatibility in Public Drinking Water Source Areas* should be referred to the Department of Water for comment. The Department will make recommendations where appropriate.




Figure 3 Land use and tenure within and surrounding the Dathagnoorara Water Reserve

4 Catchment protection strategy

4.1 Protection objectives

The objective of this plan is to protect the drinking water source in the interest of providing safe drinking water to the towns of Carnamah and Coorow, while recognising the rights of existing approved land uses to continue within the Dathagnoorara Water Reserve.

The priority classifications for the Dathagnoorara Water Reserve have been assigned to ensure consistency with this Department's current framework for public drinking water source protection. The priority classifications reflect the form of land tenure, the strategic importance of the water source, hydrogeology and land use and zoning.

4.2 Proclaimed area

The boundary of the current gazetted Dathagnoorara Water Reserve (see section 1.5.1) is proposed to be amended under the *CAWS Act 1947* to include the additional area that is formed by 300 metre Wellhead Protection Zones (WHPZ) around each of the production bores. This will include a portion of Victoria Locations 9921, 9920, 10093 and 10126. The inclusion of this additional area will allow protection of the area where contamination of the drinking water source is most likely to occur. The proposed new boundary for Dathagnoorara Water Reserve is shown in Figure 2.

4.3 Priority classifications

An explanation of the priority classification system and the detail of land use compatibility within each priority classification is provided in the Water Quality Protection Note *Land use compatibility in Public Drinking Water Source Areas*.

The area covered by the current gazetted water reserve (Crown Reserve 25198) will be classified as Priority 1 (P1) and the remainder of the proposed Dathagnoorara Water Reserve will be classified as Priority 2 (P2). Extensive agriculture is a compatible land use within the proposed P2 classification (see the Department's Water Quality Protection Note *Land use compatibility in Public Drinking Water Source Areas*). The priority classifications for Dathagnoorara Water Reserve are shown in Figure 4.

4.4 Protection Zones

The Department of Water defines Wellhead Protection Zones (WHPZ) around each production bore (500 metre radius in P1 areas and 300 metre radius in P2 and P3 areas) in which activities are to be managed to maximise protection against contamination. There are two WHPZ proposed for Dathagnoorara Water Reserve. These have a 300m radius from each bore, as shown in Figure 4.




Figure 4 Priority classifications and protection zones for Dathagnoorara Water Reserve

4.5 Land use planning

It is recognised under the *State Planning Strategy* (Western Australian Planning Commission, 1997) that the establishment of appropriate protection mechanisms in statutory land use planning processes is necessary to secure the long-term protection of drinking water sources. As outlined in the Statement of Planning Policy No.2.7 *Public Drinking Water Source Policy* (Western Australian Planning Commission, 2003) it is appropriate that the Dathagnoorara Water Reserve, protection zones and priority classifications be recognised in the Shire of Carnamah's Town Planning Scheme. Land use and activities in this water reserve should be guided by this Plan and the Water Quality Protection Note *Land Use Compatibility in Public Drinking Water Source Areas.* Any development proposals located within the Dathagnoorara Water Reserve that are inconsistent with the Department of Water's Water Quality Protection Note *Land use compatibility in Public Drinking Water Source Areas* should be referred to the Department of Water for advice and recommendations.

4.6 Best management practices

There are opportunities to significantly reduce risks to water quality by carefully considering design and management practices. The adoption of best management practices for land uses will continue to be encouraged to help protect water quality. On freehold land, the Department of Water aims to work with landowners to achieve best management practices for water quality protection through the provision of management advice, and assistance to seek funding if required.

There are guidelines available for many land uses in the form of industry codes of practice, environmental guidelines or Water Quality Protection Notes. These have been developed in consultation with stakeholders such as industry groups, producers, state government agencies and technical advisers. Examples include *Agriculture – dryland crops near sensitive water resources* and *Stockyards* (see References section). The guidelines help managers reduce the risk of their operations causing unacceptable environmental impacts. They are recommended as best practice for water quality protection.

Education and awareness (eg signage and information material) is a key mechanism for water quality protection, especially for those people visiting the area who are unfamiliar with the Dathagnoorara Water Reserve. A brochure will be produced once this Plan is endorsed, describing the Dathagnoorara Water Reserve, its location and the main threats to water quality protection. This brochure will be made available to the community and will serve to inform people in simple terms about the drinking water source and its protection.

4.7 Surveillance and By-law enforcement

The quality of public drinking water sources within country areas of the State is protected under the *CAWS Act 1947*. Declaration of these areas allows existing By-laws to be applied to protect water quality.

The Department of Water considers By-law enforcement, through on-ground surveillance of land use activities in Public Drinking Water Source Areas as an important water quality protection mechanism.

Signs are erected to educate the public and to advise of activities that are prohibited or regulated. This Plan recommends that the delegation of surveillance and By-law enforcement to the Water Corporation is continued.

4.8 Emergency response

Escape of chemicals during unforeseen incidents and the use of chemicals during emergency responses can result in water contamination. The Shire of Carnamah's Local Emergency Management Advisory Committee (LEMAC) through the Mid West-Gascoyne Emergency Management District should be familiar with the location and purpose of the Dathagnoorara Water Reserve. A locality plan should be provided to the Fire and Rescue Services headquarters for the Hazardous Materials Emergency Advisory Team (HAZMAT). The Water Corporation should have an advisory role to any HAZMAT incident in the Dathagnoorara Water Reserve.

Personnel who deal with WESTPLAN – HAZMAT (Western Australian Plan for Hazardous Materials) incidents within the area should have access to a map of the Dathagnoorara Water Reserve. These personnel should receive training to ensure an adequate understanding of the potential impacts of spills on the water resource.

4.9 Recommended protection strategies

Table 1 identifies the potential water quality risks associated with existing land uses in the Dathagnoorara Water Reserve and recommends protection strategies to minimise these risks.

Following publication of the final Dathagnoorara Water Reserve Drinking Water Source Protection Plan, an Implementation Strategy will be drawn up based on the recommendations in Table 1. It will describe timeframes and funding sources for the recommended protection strategies and identify responsible stakeholders. This is reflected in the Recommendations section of this plan. Table 1 Land use, potential water quality risks and recommended strategies

Land use / activity	Potential water quality risks		Consideration for management	Current preventative measures	Recommended protection strategies	
	Hazard	Management priority				
Broad Acre Cropping: • Wheat • Lupins • Barley • Canola	Nutrients from fertilizer application	Medium	Cereal cropping is the dominant land use in the area and all farms apply nitrogen and phosphorous with soil and plant testing often used to optimise application rates. No tillage farming is becoming more common. Depth to the water table is >90m, bores are ≥ 150m deep and aquifer is semi-confined.	 Water quality monitoring Sealed bores and fenced compounds Signage – at wellfield and Arrowsmith Groundwater Area 	Best practice management as recommended in this Department's note Agriculture – dryland crops near sensitive water resources and the Department of Agriculture and Food Farmnote series. New bores in the vicinity of the production bores should be appropriately assessed by the Department of Water	
	Pesticides	Low	Herbicides used in the area include glyphosate, simazine, atrazine and others with application at seeding and 4 to 8 weeks later if required. Pesticides used include chlorphyrifos, cypermethrin, alphamethrin and others, with application as required. Depth to the water table is >90m and aquifer is semi-confined.	 Water quality monitoring Sealed bores and fenced compounds Signage – at wellfield and Arrowsmith Groundwater Area 	Best practice management for the application of pesticides and herbicides as recommended in Statewide Policy No.2 <i>Pesticide</i> <i>use in Public Drinking Water</i> <i>Source Areas</i> and Circular No: PSC 88 Use of Herbicides in Water Catchment Areas New bores in the vicinity of the production bores should be appropriately assessed by the Department of Water.	

Land use / Potential water quality activity		er quality risks	Consideration for management	Current preventative measures	Recommended protection strategies	
	Hazard	Management priority				
Stock Grazing: • Sheep • Cattle	Nutrients from animal manure Pathogens from animal manure	Low	Sheep and cattle grazing is generally a secondary land use to cropping in the area. Stocking rates are typically low. Depth to the watertable is >90m and aquifer is semi-confined	 Water quality monitoring Sealed bores and fenced compounds Signage - at wellfield and Arrowsmith Groundwater Area 	Best practice management as recommended in the Department of Agriculture and Food's Farmnote series. New bores in the vicinity of the production bores should be appropriately assessed by the Department of Water.	
Roads	Hydrocarbons from accidents or spills	Low	Bulk haulage in the area is mainly associated with servicing farms ie haulage of grain, fertilizer and chemicals. The bores are located within 50 to 70m from Carnamah-Eneabba Rd (a major sealed road) and within 200 to 300m from Brand Mudge Road (unsealed).	 Water quality monitoring Sealed bores and fenced compounds 	Best practice management as recommended in this Department's Note <i>Roads near</i> <i>sensitive water resources.</i>	
	Pesticides from accidents or spills	Low		 Signage – at wellfield and Arrowsmith Groundwater Area LEMAC response 	Continuation of water quality monitoring, surveillance program and LEMAC response.	
	Nutrients from accidents or spills involving fertilizer	Low	Accidents and spills are rare. The Local Emergency Management Advisory Committee (LEMAC) responds to incidents.			
Mineral exploration	Hydrocarbons	Low	Exploration bores can potentially provide a pathway for contaminants to travel down into	DOIR place specific conditions on the mining tenement	Exploration drilling should be located outside of the Wellhead Protection Zones.	

Land use / activity			Consideration for management		Current preventative measures	Recommended protection strategies	
	Hazard	Management priority					
	Nutrients Pathogens	Medium Medium	the semi-confined aquifer. The tenement holder has committed to best practice management for any drilling undertaken in close proximity to the production bores.	•	with regards to the presence of a Water Reserve Water quality monitoring Sealed bores and fenced compounds	All exploration bores drilled within close proximity of the production bores should be backfilled and sealed in an approved manner.	
Intensive Agriculture (potential)	Nutrients Pesticides Hydrocarbons	Medium	Under the current land use zoning there is the potential for intensification of agricultural activities within and surrounding the proposed Water Reserve. Depth to groundwater is >90m, bores are ≥ 150m deep and aquifer is semi-confined.	•	Water quality monitoring Land use planning approvals process	Relevant local government authorities to refer development proposals that are inconsistent with the Department's Land Use Compatibility Table to the Department of Water for advice and recommendations. New bores in the vicinity of the production bores should be appropriately assessed by the Department of Water.	

5 Recommendations

- 1 Implement the recommended protection strategies as detailed in *Table 1: Land* use, potential water quality risks and recommended strategies of this Plan (Applicable stakeholders).
- 2 The boundary of the Dathagnoorara Water Reserve should be amended under the *Country Areas Water Supply Act 1947 (Department of Water).*
- 3 Prepare an implementation strategy for this Plan describing responsible stakeholders, timeframes and funding sources for the recommended protection strategies (*Department of Water*).
- 4 The Shire of Carnamah Town Planning Scheme should incorporate this Plan and reflect the identified Dathagnoorara Water Reserve boundary, the Priority 1 and Priority 2 classifications and the protection zones (*Shire of Carnamah*).
- 5 All development proposals within the Dathagnoorara Water Reserve that are inconsistent with the Water Quality Protection Note *Land use compatibility in Public Drinking Water Source Areas* or the Statement of Planning Policy No. 2.7 *Public Drinking Water Source Policy* should be referred to the Department of Water for advice and recommendations (*Department for Planning and Infrastructure, Shire of Carnamah*).
- 6 Applications to construct a bore and/or extract groundwater in close proximity to the production bores within the Dathagnoorara Water Reserve should be assessed to ensure that the bores are appropriately located. Best management practices should be recommended for the maintenance and construction of the bores to prevent potential contamination or reduction in water availability to the public drinking water source bores (*Department of Water*).
- 7 Exploration bores drilled in close proximity to the production bores must be backfilled and sealed in an approved manner as soon as possible (*Mining tenement holder, Department of Water*).
- 8 Any new exploration bores should be located outside of the Wellhead Protection Zones. Proposals to undertake exploration drilling within the Wellhead Protection Zones should be forwarded to the Department of Water for advice and recommendations (*Mining tenement holder, Department of Industry and Resources*).
- 9 Incidents covered by WESTPLAN HAZMAT in the Dathagnoorara Water Reserve should be addressed through the following:
 - the Mid West-Gascoyne LEMAC are familiar with the location and purpose of the Dathagnoorara Water Reserve;
 - the locality plan for the Dathagnoorara Water Reserve is provided to the Fire and Rescue headquarters for the HAZMAT Emergency Advisory Team;
 - the Water Corporation provides an advisory role during incidents in the Dathagnoorara Water Reserve; and

 personnel dealing with WESTPLAN – HAZMAT incidents in the area have ready access to a locality map of the Dathagnoorara Water Reserve and training to understand the potential impacts of spills on drinking water quality

(Department of Water, Water Corporation).

- 10 The current surveillance program should be maintained to identify any incompatible land uses or potential threats to the Dathagnoorara Water Reserve (*Water Corporation*).
- 11 Signs located along the boundary of the Dathagnoorara Water Reserve should be maintained to define its location and promote awareness of the need to protect drinking water quality. Signs should include an emergency contact telephone number (*Water Corporation*).
- 12 A review of this Plan should be undertaken after five years in consultation with the Shire of Carnamah and the affected private property owners (*Department of Water*).

Appendices

Appendix A - Summary of Submissions

The following table is a summary of the submissions received from the 2007 release of the Draft Plan, the current status of that issue, and how they have been addressed in this current Plan.

Summary of submissions from the 2007 release of the Draft Plan

Issue raised	Current status	Response in Plan/Comment
Recommendation 10 should include the statement 'in consultation with the Shire of Carnamah and affected private land owners'.	Any review of the plan will include consultation with all relevant stakeholders.	Statement added to Recommendation 10.
Cost of amendment of the Shire of Carnamah's Town Planning Scheme – should be borne by the Department of Water.	Shire of Carnamah are not required to amend their TPS immediately, however, when the TPS is next reviewed the Dathagnoorara Water Reserve should be included in it as a special control area.	No change in plan.
Potential for purchasing private property within the Water Reserve rather than imposing controls on land owners.	Purchase of privately owned land is an option, however, is only undertaken on a willing seller basis. It will also depend upon the Department of Water's resources and priorities for purchasing land. The land owners have not indicated that they wish to sell their land at this stage.	No change in plan.
Concern that there may be an adverse effect on borrowing potential for private landowners with land within the Water Reserve.	Due to the very small proportion of each block of land covered by the Water Reserve, it is unlikely that the extension of the Water Reserve boundary will have a significant impact on the land owners borrowing potential.	No change in plan.
Mining tenement holder of M70/2759 (part of which lies over Dathagnoorara Water Reserve) is undertaking a mineral exploration program on this tenement, which may include drilling within the Water Reserve.	Mining tenement holder has committed to a number of management approaches to ensure there is no increased risk to the drinking water source.	Recommendations 7 and 8 added to the plan.

Appendix B- Water quality

The information provided in this appendix was developed by the Water Corporation's Water Quality Branch.

The Water Corporation has monitored the raw (source) water quality from Dathagnoorara borefield in accordance with the Australian Drinking Water Guidelines (ADWG) and interpretations agreed to with the Department of Health. The raw water is regularly monitored for:

- a. aesthetic related characteristics (non-health related)
- b. health related characteristics including:
 - health related chemicals; and
 - microbiological contaminants.

Following is data representative of the quality of raw water from the Dathagnoorara borefield. In the absence of specific guidelines for raw water quality, the results have been compared with the ADWG values set for drinking water, which defines the quality requirements at the customers tap. Results that exceed the ADWG have been shaded to give an indication of potential raw water quality issues associated with this source.

It is important to appreciate that the raw water data presented does not represent the quality of drinking water distributed to the public. Barriers such as storage and water treatment, to name a few, exist downstream of the raw water to ensure it meets the requirements of the ADWG. For more information on the quality of drinking water supplied to Carnamah/Coorow Scheme refer to the most recent Water Corporation Drinking Water Quality Annual Report at

<www.watercorporation.com.au/W/waterquality_annualreport.cfm?uid=2377-9937-9579-7091>.

Aesthetic related characteristics

Aesthetic water quality analyses for raw water from Dathagnoorara borefield are summarised in the following table.

The values are taken from ongoing monitoring for the period January 2002 to May 2007. All values are in milligrams per litre (mg/L) unless stated otherwise. Any water quality parameters that have been detected are reported, those that have on occasion exceeded the ADWG are shaded.

The raw water from the Dathagnoorara wellfield complies with most ADWG aesthetic guidelines. Turbidity, chloride, iron, sodium and pH concentrations have been measured at levels exceeding the guidelines. Salinity (TFSS) has consistently been measured at levels above the ADWG aesthetic level (500mg/L), however it is stated in the ADWG that a level between 500-1000mg/L is acceptable based on taste.

Parameter	Units	ADWG Aesthetic	Dathagnoorara Borefield Raw Source SP		
		Guideline Value*	Range	Median	
Aluminium unfiltered	mg/L	NA	<0.008 - 0.032	<0.008	
Chloride [†]	mg/L	250	420	420	
Conductivity at 25°C	mS/m	NA	140 - 165	150	
Hardness as CaCO₃ [†]	mg/L	200	115 - 123	119	
Iron unfiltered	mg/L	0.3	<0.003 - 0.34	0.008	
Manganese unfiltered	mg/L	0.1	0.03 - 0.05	0.04	
рН	-	6.5 - 8.5	5.04 - 5.36	5.16	
Sodium [†]	mg/L	180	245 - 255	250	
Sulphate [†]	mg/L	250	48 - 50	49	
TFSS [†]	mg/L	500	786 - 804	795	
Turbidity	NTU	5	<0.1 - 6.4	<0.1	

Aesthetic related	detections	for Dathagnoorara	borefield
ACSTRUCTURE		joi Duchuznooruru	borcjicia

* An aesthetic guideline value is the concentration or measure of a water quality characteristic that is associated with good quality water.

[†]Water quality data observed from 2 sampling occasions

Health related characteristics

Health Parameters

Raw water from Dathagnoorara borefield is analysed for health related chemicals including inorganics, heavy metals, industrial hydrocarbons and pesticides. Health related water quality parameters that have been measured at detectable levels in the source between January 2002 and May 2007 are summarised in the following table. All health related water quality parameters detected were within ADWG levels and pose no health concern.

Parameter	Units	ADWG Health Guideline	Dathagnoorara Borefield Raw Source SP		
		Value*	Range	Median	
Barium [†]	mg/L	0.7	0.075	0.075	
Boron [†]	mg/L	4	0.08 - 0.12	0.1	
Manganese unfiltered	mg/L	0.5	0.03 - 0.05	0.04	
Nitrate as nitrogen [†]	mg/L	11.29	<0.002 - 0.1	<0.051	
Nitrite as nitrogen [†]	mg/L	0.91	0.003 - 0.004	0.0035	
Nitrite plus nitrate as N [†]	mg/L	11.29	0.11 - 0.17	0.14	
Sulphate [†]	mg/L	500	48 - 50	49	

Health related detections for Dathagnoorara borefield

* A health guideline value is the concentration or measure of a water quality characteristic that, based on present knowledge, does not result in any significant risk to the health of the consumer over a lifetime of consumption (NHRMC & ARMCANZ, 1996).

[†]Water quality data observed from 3 or less sampling occasions

Microbiological contaminants

Microbiological testing of raw water samples from Dathagnoorara borefield is currently conducted on a monthly basis. *Escherichia* coli counts are used as an indicator of the degree of recent faecal contamination of the raw water from warmblooded animals. A detection of *Escherichia coli* in raw water abstracted from any bore may indicate possible contamination of faecal material through ingress in the bore, or recharge through to the aquifer (depending on aquifer bore type).

During the reviewed period of January 2002 to May 2007, no positive *Escherichia coli* counts were recorded in any samples collected from the borefield. This is indicative of minimal contamination of the groundwater from faecal sources.

Appendix C- Photographs



Photo 1: Water Corporation Production Bore 2/86 and vegetated reserve



Photo 2: Typical rural land use within the region

Glossary

Abstraction	The pumping of groundwater from an aquifer.		
ADWG	The Australian Drinking Water Guidelines, outlining guideline criteria for the quality of drinking water in Australia.		
Aesthetic guideline	NHMRC guideline level ascribed to acceptable aesthetic qualities of drinking water such as taste, smell, colour and temperature.		
AHD	Australian Height Datum is the height of land in metres above mean sea level. For example this is +0.026 m at Fremantle.		
Allocation	The quantity of water permitted to be abstracted by a licence, usually specified in kilolitres per year (kL/a).		
ANZECC	Australian and New Zealand Environment Conservation Council.		
Aquifer	A geological formation or group of formations able to receive, store and transmit significant quantities of water.		
ARMCANZ	Agriculture and Resource Management Council of Australia and New Zealand.		
Bore	A narrow, lined hole, also known as a well, drilled to monitor or draw groundwater.		
Borefield	A group of bores to monitor or withdraw groundwater.		
Catchment	The area of land which intercepts rainfall and contributes the collected water to surface water (streams, rivers, wetlands) or groundwater.		
CFU	Colony forming units is a measure of pathogen contamination in water.		
Confined aquifer	An aquifer that is confined between non-porous rock formations (such as shale and siltstone) and therefore contains water under pressure.		
HAZMAT	Hazardous Materials		
Hydrogeology	The study of groundwater, especially relating to the distribution of aquifers, groundwater flow and groundwater quality.		
kL	Kilolitres (1000 litres)		
km	Kilometres (1000 metres)		
LEMAC	Local Emergency Management Advisory Committee		

m	Metres		
mg/L	Milligrams per litre (0.001 grams per litre)		
mm	Millimetres		
MPN	Most probable number, a measure of microbiological contamination		
mSv	Millisieverts, a measure of annual radiological dose		
NHRMC	National Health and Medical Research Council.		
NTU	Nephelometric turbidity units are a measure of turbidity in water.		
Nutrients	Minerals dissolved in water, particularly inorganic compounds of nitrogen (nitrate and ammonia) and phosphorous (phosphate) which provide nutrition (food) for plant growth. Total nutrient levels include the inorganic forms of an element plus any bound in organic molecules.		
Pesticides	Collective name for a variety of insecticides, fungicides, herbicides, algicides, fumigants and rodenticides used to kill organisms.		
Public Drinking Water Source Area (PDWSA)	Includes all underground water pollution control areas, catchment areas and water reserves constituted under the <i>Metropolitan Water Supply Sewerage and Drainage Act 1909</i> and the <i>Country Areas Water Supply Act 1947</i> .		
Recharge	Water infiltrating to replenish an aquifer.		
Recharge area	An area through which water from a groundwater catchment percolates to replenish (recharge) an aquifer. An unconfined aquifer is recharged by rainfall throughout it's distribution. Confined aquifers are recharged in specific areas where water leaks from overlying aquifers, or where the aquifer rises to meet the surface.		
Scheme supply	Water diverted from a source or sources by a water authority of private company and supplied via a distribution network to customers for urban, industrial or irrigation use.		
Semi-confined aquifer	A semi-confined or leaky aquifer is saturated and bounded above by a semi-permeable layer and below by a layer that is either impermeable or semi-permeable.		
TCU	True colour units, a measurement of colour in water.		
TDS	Total dissolved salts, a measurement of ions in solution, such as salts in water.		

Treatment	Application of techniques such as settlement, filtration and chlorination to render water suitable for specific purposes including drinking and discharge to the environment.		
Unconfined aquifer	An aquifer in which the upper surface of water is lower than the top of the aquifer itself. The upper surface of the groundwater within the aquifer is called the watertable.		
Water quality	The physical, chemical and biological measures of water.		
Water Reserve	An area proclaimed under the <i>Country Areas Water Supply Act 1947</i> of the <i>Metropolitan Water Supply Sewerage and Drainage Act 1909</i> for the purposes of protecting a drinking water supply.		
Watertable	The upper saturated level of the unconfined groundwater.		
Wellfield	A group of bores to monitor or withdraw groundwater.		
Wellhead	The top of a well (or bore) used to draw groundwater. A wellhead protection zone (WHPZ) is usually declared around wellheads in drinking water areas to protect the water source from contamination.		
WESTPLAN HAZMAT	Western Australian Plan for Hazardous Materials.		

References and Recommended Reading

- Agriculture and Resource Management Council of Australia and New Zealand (ARMCANZ) & Australian and New Zealand Environment and Conservation Council (ANZECC) 1996, *Draft Rural Land Uses and Water Quality – A Community Resource Document*, ARMCANZ & ANZECC, Canberra.
- Bureau of Meteorology 2007, *Climate statistics for Australian Locations: Summary statistics Carnamah*, Bureau of Meteorology, available http://www.bom.gov.au/climate/averages/tables/cw_008025.shtml.
- Department of Agriculture and Food Various dates, *Farmnotes –Various titles,* Department of Agriculture and Food, Perth, available <www.agric.wa.gov.au/servlet/page?_pageid=449&_dad=portal30&_schema=P ORTAL30> > Publications > Farmnotes.
- Department of Health 2007, *Use of Herbicides in Water Catchment Areas*, Circular No: PSC 88, February 2007, Department of Health, Perth.
- Department of Water Various dates, *Water Quality Protection Notes Various titles*, Department of Water, Perth, available <www.water.wa.gov.au> > Water Quality > Publications > Water Quality Protection Notes.
- Government of Western Australia 1914, *Rights in Water and Irrigation Act*, Reprinted under the *Reprints Act 1984* as at 4 July 2006, available .
- Government of Western Australia 1947, *Country Areas Water Supply Act*, reprinted under the Reprints Act 1984 as at 7 April 2006, available http://www.slp.wa.gov.au/statutes/swans.nsf>.
- National Health and Medical Research Council (NHMRC) & Natural Resource Management Ministerial Council (NRMMC) 2004, National Water Quality Management Strategy: Australian Drinking Water Guidelines, Australian Government, Canberra, available <www.nhmrc.gov.au/publications/synopses/eh19syn.htm>.

NHMRC & NRMMC 2004b, Water Made Clear – A consumer guide to accompany the Australian Drinking Water Guidelines 2004, Australian Government, Canberra.

National Minimum Bore Specifications Committee 2003, *Minimum construction requirements for water bores in Australia*, 2nd edn, Land and Water Biodiversity Committee, Queensland, available

<http://portal.water.wa.gov.au/portal/page/portal/LicensingWaterIndustryServices /Licensing/Publications/Content/MCRWBA.pdf>

- Water and Rivers Commission 2000, Statewide Policy No.2 *Pesticide Use in Public Drinking Water Source Areas*, Water and Rivers Commission, Perth, available <www.water.wa.gov.au> > Policies.
- Water and Rivers Commission 2002, Interim Sub-Regional Allocation Strategy Managing the Water Resources of the Arrowsmith Groundwater Area WA, Water and Rivers Commission, Perth.
- Water Authority 1995, Arrowsmith Groundwater Area Management Plan Report No. WG153, Water Corporation, Perth.
- Water Corporation 2004, Dathagnoorara Water Reserve Drinking Water Source Protection Assessment, Water Corporation, Perth.
- Water Corporation 2006, *SG097 Source Protection Operations Manual*, Water Corporation, Perth.
- Western Australian Planning Commission 1997, *State Planning Strategy*, State Government of Western Australia, Perth, available www.wapc.wa.gov.au/Publications/52.aspx.
- Western Australian Planning Commission 2003, Statement of Planning Policy No. 2.7 Public Drinking Water Source Policy, Government Gazette WA, 10 June 2003, pp. 2077-82, Government of Western Australia, Perth, available <www.wapc.wa.gov.au/Publications/149.aspx>.
- Western Australian Planning Commission 2006a, *Shire of Carnamah Town Planning Scheme No. 1,* Department for Planning and Infrastructure, Perth, available www.wapc.wa.gov.au/Planning/Schemes/Local Planning Schemes.
- Western Australian Planning Commission 2006b, Statement of Planning Policy No.
 2.9 Water resources, Government Gazette WA, 19 December 2006, pp 570722, Government of Wester Australia, Perth, available
 <www.wapc.wa.gov.au/Publications/1281.aspx>.
- World Health Organisation 1996, Guidelines for Drinking-Water Quality Volume 2 Health Criteria and Other Supporting Information, 2nd edn, World Health Organisation, Geneva, available <www.who.int/water_sanitation_health/dwq/en/2edvol2p1.pdf>.

Contributors

This report was prepared by:

Contribution	Personnel	Title	Organisation
Supervision	Tony Laws	Branch Manager, Water Source Protection	Department of Water
	Nigel Mantle	A/Program Manager, Water Source Protection	Department of Water
Report Preparation	Alana Thorpe	Environmental Officer, Water Source Protection	Department of Water
Project Liaison	Luke Richards	Regional Hydrogeologist, Midwest Gascoyne Region	Department of Water
	Tran Huynh	Catchment Co-ordinator, Mid West-Gascoyne Region	Water Corporation
Drafting	Melanie Webb	GIS Officer	Department of Water
Photographs	Alana Thorpe	Environmental Officer	Department of Water

